

TPLO (Tibial Plateau Levelling Osteotomy) for Cranial Cruciate Ligament Injury

Your vet has arranged for me to see your dog because they are concerned that there is an injury to the *cranial cruciate ligament* in their knee joint. This is a fairly common injury in dogs. You may have heard of cruciate ligament injuries in people – it's a common knee injury in footballers and other athletes.

Although both dogs and people injure this ligament, the cause of the problem appears to be quite different. In people, a structurally normal ligament is snapped by excessive force. In dogs it appears that the ligament gradually deteriorates over time, for reasons that we don't fully understand. This means that very often there is a period during which the knee is sore even though the ligament hasn't snapped, and when the ligament does give way, it may be during completely normal exercise without there being any accident/fall.

Although our understanding of the underlying reason for the ligament to degenerate is poor, what we know very clearly is that in dogs that are heavier than about 10-15kg, surgical treatment is usually required. Treatment without surgery is usually unsatisfactory and leads to a chronic lameness. Surgery may also be recommended in some smaller dogs if they have not recovered by themselves or sometimes where there are underlying problems with the bone which have contributed to the problem and need to be corrected.

The aim of surgery is to stabilise the knee. Damage to the cranial cruciate ligament results in instability of the knee in a forward/backward direction. When your dog is attempting to bear weight on the affected leg, the knee is giving way so they cannot push off that leg with any confidence. Historically, surgery consisted of a variety of different ways to replace the ligament, with wire/nylon/carbon fibre or with tissue taken from elsewhere in the body. Unfortunately many of these procedures were unsatisfactory, essentially because nothing does the job of a ligament as well as a ligament. Many of the materials we used to replace the ligament would stretch/break too quickly and the surgery would be unsuccessful.

About 15-20 years ago, a new approach to the problem was introduced, and has subsequently become the accepted gold-standard treatment for cruciate ligament problems in dogs. This approach consists of altering the way the knee works to make it stable without the ligament. This is achieved by making a cut in the top of the shinbone, rotating a piece of bone slightly, and fixing it back in place with a plate and screws. The benefits of this procedure are that once the bone has healed in the new position there is nothing to stretch or break, and the knee is stable for life. Since the procedure was introduced, many hundreds of thousands of dogs worldwide have been treated in this way and there is a large body of evidence demonstrating that it is effective. The downside of the procedure is that it is more technically demanding and requires a lot of specialised equipment, meaning that many general vets do not perform it and need to refer dogs to a specialist orthopaedic surgeon for this treatment.

The Operation

Your practice will make arrangements with you to drop your pet off on the day of the surgery. Please don't give breakfast on the morning of the procedure. If they are receiving pain relief, they can have their pain relief the night before the operation, but if you normally give it in the morning then please don't give it on the day. Please let the nurse/vet admitting your dog know when they last had pain relief, and also if you will need a further supply of any medication. I will probably not meet you in person on the day of the procedure, but I will speak to you by phone if possible. If you have any questions based on the information in this sheet then please feel free to ask me then. When I arrive at the practice I will examine your dog and confirm the diagnosis and plan for surgery. If for any reason we need to alter the plan I will contact you to discuss it.

Your dog will be mildly sedated to minimise stress, and then anaesthetised. We may need some further x-rays prior to surgery to make measurements. Your dog's leg will be prepared for surgery. The leg will be shaved from hip to toe to give us a large clean zone to minimise the risk of infection.

The first part of the operation is to open the knee joint and remove the damaged remnants of the ligament. We also need to check structures called the menisci ("cartilages") which can also be damaged/torn. If the menisci are damaged then the damaged parts will be removed.

Once the joint has been inspected the procedure to cut the bone, rotate the piece and reattach it with plate and screws will be performed. Then the incision will be closed with stitches and/or staples. The leg will not be bandaged, unless there is a concern about swelling.

The whole procedure usually takes about 60-75 minutes. X-rays will be taken after the procedure to ensure that everything is in a good position. Your dog will probably be anaesthetised for about 2 hours in total because of the preparation beforehand and for the x-rays and cleaning/dressing afterwards. I will call you with an update at the end of the operation.

In most cases your dog will be discharged from the practice the evening of the procedure. You will receive a printed discharge note from me which will give details of the procedure, medication and instructions for exercise. Your dog will receive an injection of painkiller immediately before going home and you will also generally have some further pain relief to give at bedtime. Nonetheless, on the night after a major operation like this you may find your pet is restless or agitated. You should make sure that you know how to contact your practice's out-of-hours service and also where you would need to go if your dog needed to be checked over, bearing in mind that this may not be at your usual practice premises.

EXERCISE & RECOVERY

Before surgery, please obtain a crate or cage for your dog to be confined. This should be about twice the size they need to lie down comfortably. Your dog will be strictly rested for the first 6 weeks after surgery. More detailed instructions will be included in the discharge note.

Risks of Surgery

This is a procedure with relatively low risks, and it is a procedure that for us is very routine. However as you would expect for major surgery there are some risks.

1. Infection – Infection is a small risk of all surgery. We will take all the same precautions to avoid infection as I would take if I were performing the surgery at a specialist hospital (drapes, gowns, gloves etc). Your dog will receive antibiotics during surgery and will also have a short course of antibiotics after surgery. You should monitor the wound after surgery for any excessive redness, swelling or discharge. If an infection does develop it will generally resolve with further medical treatment. Very occasionally the area may need to be flushed. In very rare cases an infection may result in a poor outcome or may mean that the plate and screws need to be removed.
2. “Late Meniscal Injury” – if the cartilages in the knee are normal at the time of surgery, we will leave them in place, as they are important for knee function. Occasionally a cartilage tear will occur later on and result in a recurrence of lameness. A surgical procedure to treat the cartilage tear may be required.
3. Osteoarthritis – Some degree of “wear and tear” in the joint is inevitable following the injury and no surgical procedure has ever been demonstrated to stop this. However, surgical treatment now is the best way to minimise this, and many dogs will not be troubled by arthritis for many years.
4. Other leg affected – because this appears to be a “condition” rather than an injury in dogs, the other knee is very often affected within 1-2 years and the same treatment required. There is not a great deal that can be done to prevent this other than keeping your dog at a sensible weight.

There is a small risk associated with a general anaesthetic. Your vet may discuss with you whether you would like your dog to have a blood test prior to surgery to ensure that there are no underlying health problems that we were unaware of.

Thank you for entrusting the care of your pet to us, and I look forward to speaking to you on the day of the procedure.

Dr Richard Coe MA VetMB CertSAS MRCVS
Diplomate of the European College of Veterinary Surgeons (DipECVS)
RCVS & European Registered Specialist in Surgery

Dr Nick Goldfinch BVMS PhD PgC(Surg) GPCerts(SAS, SAM) MRCVS
RCVS Advanced Practitioner in Small Animal Surgery